

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 9/20</p>	<p>A2</p>	<p>(11) International Publication Number: WO 98/20860 (43) International Publication Date: 22 May 1998 (22.05.98)</p>
<p>(21) International Application Number: PCT/US97/20217 (22) International Filing Date: 13 November 1997 (13.11.97) (30) Priority Data: 08/746,733 15 November 1996 (15.11.96) US (71) Applicant: FMC CORPORATION [US/US]; 1735 Market Street, Philadelphia, PA 19103 (US). (72) Inventors: BUBNIS, William, A.; 5904 Ravens Crest Drive, Plainsboro, NJ 08536 (US). REILLY, William, J., Jr.; 409 Byerly Drive, New Hope, PA 18938 (US). (74) Agent: CALDWELL, John; Woodcock Washburn Kurtz Mackiewicz & Norris, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>
<p>(54) Title: HYDROCOLLOID DELIVERY SYSTEM</p> <p>(57) Abstract</p> <p>A chewable delivery system stable in acidic, neutral or alkaline systems and having good mouthfeel is disclosed.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HYDROCOLLOID DELIVERY SYSTEM

5 Field of the Invention

This invention relates to a delivery system for pharmacologically active materials. More particularly, this invention relates to a chewable delivery system which has good mouthfeel and which is capable of being utilized in an acidic, neutral or alkaline system.

10

Background of the Invention

When medication is prescribed and needs to be taken orally at regular intervals, the medication many times is distasteful, for example, cholestyramine. The distasteful nature of the medication often leads to failure of the patient to take
15 the medication in the recommended dosages and at the scheduled intervals.

In order to overcome the reluctance of the patient to take the medication in the prescribed dosages and at the scheduled intervals, numerous delivery systems have been devised to make the taking of the medication more palatable to the patient.

20

One such delivery system is disclosed in US Patent 4,950,689. The '689 patent discloses a delivery system made from a pectin gel and an ingestible insoluble solid added to the pectin gel product in order to provide sufficient internal matrix strength to the gel. Further, the '689 patent teaches that the pectin
25 gel, with the insoluble addition, can only be used in acidic pH so that the pH will be within the gelling range of the pectin to provide adequate gellation.

US Patent 4,695,463 is directed to a delivery system for active ingredients which comprises an insolubilized active ingredient and a crosslinked alginate or carrageenenate matrix which entraps an insolubilized active ingredient. The '463
30 patent teaches that a monovalent carrageenenate or alginate is dissolved in water with the active material and that the monovalent metal cations are then displaced

by multivalent cations, i.e., calcium to form a crosslinked matrix which will then precipitate. The precipitate is then recovered and dried to form the delivery system.

5 European Patent Publication 0 075 443 is directed to a delivery system for an antacid which is a chewable composition and which is prepared from a light textured frappe.

Canadian Patent Application 2044043 is directed to a delivery system for a
10 drug and other materials, wherein the delivery system is in the form of a film and which may contain various gelling agents.

None of the aforescribed references teach or suggest a chewable delivery system for a pharmacologically active material, which delivery system is a low
15 moisture hydrocolloid containing system. Further, none of the references teaches a delivery system which provides a smooth creamy method of delivering pharmacological ingredients which otherwise have an unpleasant taste or texture so that an otherwise unpleasant pharmacological active material is ingested by the patient at the scheduled intervals in the prescribed dosage.

20

Summary of the Invention

Broadly, this invention contemplates a chewable delivery system for a pharmacologically active material comprising sweetener, a hydrocolloid and water, said delivery system comprising from about 50% to about 83% of solids.

25

Unless indicated otherwise, all percentages given herein are percentages by weight based on the weight of the entire composition.

Detailed Description

30 The term pharmacological, as used in the specification and claims means a material useful to treat a medical condition or having nutritive value.

A delivery system for a pharmacologically active material would be prepared by heating deionized water with stirring and adding a hydrocolloid such as carrageenan to the hot water and dissolving it therein. After the hydrocolloid is fully dissolved, sucrose is added to the aqueous solution which is stirred and heated until the sucrose is fully dissolved. A second syrup solution is prepared by placing corn syrup and water in a stainless steel beaker, with overhead stirring, and heating it in a boiling water bath. Thereafter, sucrose is added to the second solution with stirring until the sucrose dissolves completely. A portion of the carrageenan-containing solution is then added to the second solution with stirring until the mixture is uniform. Thereafter, a pharmacologically active material such as calcium carbonate is added and mixing is continued until it is uniformly dispersed. The beaker which contains the calcium carbonate is then heated to evaporate water until a predetermined weight is achieved. Thereafter, artificial flavoring may be added and the mixture may then be poured into suitable molds. The mold is then allowed to cool and the contents solidify. After cooling, the resultant gel is then ready to be ingested without the gritty chalky texture of the pharmacological component (calcium carbonate).

In preparing the chewable delivery system of this invention, it is important that the solids content of the delivery system be from about 50% to about 83%. In a preferred embodiment, the solids content of the system should be from about 78% to about 83%. When the solids content is between about 78% to about 83%, then a preservative may not be necessary for the composition because the delivery system may be stored without supporting microbial growth thereon.

When the solids content of the delivery system is between about 50% to up to about 78%, it is desirable to include an anionic or neutral preservative such as sodium benzoate, potassium sorbate, or a paraben in an amount of from about 0.02% to about 0.5%.

When preparing the delivery system, the amount of solids present is calculated and any excess water is removed in any satisfactory manner such as by boiling.

When removing the excess water, attention should be given to the

pharmacologically active material which is included in the delivery system. For

5 example, acetaminophen may be partially or completely decomposed by heat and moisture. Therefore, care should be taken to incorporate such pharmacologically active materials into the molten formulation as rapidly and as late as possible to enable the delivery system to be useful for the heat sensitive active ingredients.

10 When carrageenan is used as the hydrocolloid, the carrageenan which is used is sulfated to the extent of about 18% to about 35% based on the weight of the molecule.

The amount of hydrocolloid, such as carrageenan, which is used may vary
15 widely from about 2% to about 5.5%. Although an amount in excess of 5.5% may be used, there is no advantage to using such a large amount.

The delivery system may also include one or more additional hydrocolloids in a total amount of from about 0.5% to about 2%. Any suitable additional
20 hydrocolloid which is compatible with the delivery system and the pharmacologically active material which is incorporated and which is stable at the pH ranges set forth herein may be used. Among such additional hydrocolloids may be mentioned locust bean gum, a polysaccharide sulfated to an extent greater than 35% based on the weight of the molecule, konjac mannan, a starch, guar gum
25 and the like.

The pH of this delivery system may vary widely from about 2 to about 11. The fact that the pH of the delivery system may vary as widely as set forth, renders the delivery system useable with a wider range of pharmacologically active
30 materials than other delivery systems. Any suitable sweetener may be used in the delivery system. The sweetener used in the delivery system of this invention may

be a natural sweetener, such as sucrose, fructose, xylose, ribose, glucose, mannose, galactose, corn syrup, and mixtures thereof and the like.

If desired, an artificial sweetener may also be used such as saccharin salts, aspartame, cyclamate salts, dipeptide based sweeteners, talen, dihydrochalcone, hydrogenated starch hydrolysate, sugar alcohols, and the like and mixtures thereof.

The amount of sweetener used will vary widely and is determined by what is calculated to be the desired solids content of the final composition. Thus, one need only calculate the amount of the solids being added to the water and the amount of water which must be removed to give the desired solids content.

If desired, a flavoring agent and a coloring agent may also be incorporated in the delivery system and may be selected from any of those deemed suitable for use in food and drug applications providing they are compatible with the remainder of the delivery system. The amount of flavoring agent used may be chosen from representative flavoring agents such as spearmint oil, cinnamon oil, oil of wintergreen, peppermint oils, fruit flavors such as citrus oil including lemon, orange, grape, lime and grapefruit and fruit essences including apple, strawberry, cherry, pineapple and the like. In addition, powdered flavorings may also be used.

The amount of flavoring agent employed is generally a matter of preference. However, a flavoring agent may be present in an amount of from about 0.05% to about 3%.

If a colorant is used, a pigment, such as titanium dioxide, may be used in an amount of up to about 1% and preferably up to about 0.6%. Any suitable colorant may be used.

The delivery system of this invention may be used to deliver a wide variety of pharmacological materials such as antacids and many other useful drugs.

Examples of useful pharmacologically active materials include: mineral supplements, analgesics, antipyretics, ion exchange resins, appetite suppressants, vitamins, anti-inflammatory substances, coronary dilators, cerebral dilators, peripheral vasodilators, anti-infectives, psychotropics, antimanics, stimulants, antihistamines, laxatives, decongestants, gastro-intestinal sedatives, antidiarrheal preparations, anti-anginal drugs, vasodilators, antiarrhythmics, anti-hypertensive drugs, vasoconstrictors and migraine treatments, antibiotics, tranquilizers, antipsychotics, antitumor drugs, anticoagulants and antithrombotic drugs, hypnotics, sedatives, antiemetics, anti-nauseants, anticonvulsants, neuromuscular drugs, hyper- and hypoglycemic agents, thyroid and antithyroid preparations, diuretics, antispasmodics, uterine relaxants, mineral and nutritional additives, antiobesity drugs, anabolic drugs, erythropoietic drugs, antiasthmatics, expectorants, cough suppressants, mucolytics, anti-uricemic drugs and mixtures thereof.

In order to more fully illustrate the nature of this invention and the manner of practicing the same, the following examples are presented.

Example 1

Chewable Calcium Carbonate Antacid Carrageenan-based Confection

Deionized water (460 grams) was placed in a 2 liter, stainless steel beaker that was heated by a boiling water bath. The water was stirred with an overhead stirrer, and 19.85 grams of carrageenan (Gelcarin® GP-911, FMC Corporation, Philadelphia, PA 19103) was dissolved in the hot water. When the carrageenan was fully dissolved, 20.15 grams of sucrose was added to the aqueous solution. Stirring and heating continued until the sucrose was fully dissolved. Simultaneously, a syrup solution was prepared by placing 409.6 grams of corn syrup (42/43 DE, 80% solids by weight) and 54.4 grams of water in a 2 liter stainless steel beaker that was stirred with an overhead stirrer and heated in the boiling water bath. Next, 255.7 grams of sucrose was added to the beaker and stirring was continued until the sucrose had dissolved completely. A portion (280.3

grams) of the aqueous carrageenan solution was added to the syrup solution. After the mixture had become uniform, 100 grams of calcium carbonate (USP/NF) was added to it, and mixing was continued until it was uniformly dispersed. The beaker was transferred to a hot plate and heated to evaporate water until a predetermined weight of 842.362 grams was achieved. A solution of 2.5 grams of artificial wintergreen flavor (59.483, Firmenich, Inc.) dissolved in 20 grams of water was added and mixed thoroughly after which the mixture was poured into tubular molds. The mixture that was poured into the molds comprised 79.5% solids by weight. The molds were made from copper tubing which was capped at one end. For filling, the mold was held in a vertical position with the capped end at the bottom. After a mold was filled with hot solution, it was allowed to cool, the contents solidifying as it cooled. By removing the cap from the closed end of the mold, it was possible to push the solid, gelled product from the mold. Immediately, the gelled product was rolled in powdered starch to overcome any residual stickiness. It was calculated that 4.431 grams of gelled product contained 500 mg of calcium carbonate. Portions of the product weighing approximately 4.4 grams were sliced from the cylinder of product and chewed. The gel had a soft, chewy, creamy texture with the flavor of wintergreen predominating. The inherent gritty, chalky texture of the calcium carbonate was completely eliminated by the gel, which caused no toothpacking or stickiness.

Example 2

Chewable Calcium Carbonate Antacid Carrageenan-based Confection with Lower Solids Content

The method of Example 1 was followed exactly except that the water was not evaporated to the same extent, the flavor was changed to punch flavor (586.323/AP, Firmenich, Inc.), and FD&C Red #40 food coloring (0.20 gram) was added to the formulation. The solids content of this mixture was 66.0%. The solidified gel was more brittle than the gel of Example 1, but the organoleptic properties of both gels were very similar. After a storage period, syneresis of water and bacterial growth in the gel occurred, indicating that the solids, i.e., sugar,

content was too low to prevent these undesirable occurrences. No preservative had been added to this formulation to prevent the microbial growth.

Example 3

5 Chewable Calcium Carbonate Antacid Carrageenan-based Confection with
Increased Calcium Carbonate Content

10 The method of Example 1 was followed exactly except that 200 grams of calcium carbonate was incorporated into the formulation and butterscotch flavoring was added in place of wintergreen flavoring. Before pouring the mixture into the molds, it was determined that the solids content of this formulation was 82.2% by weight. Each 5.08 gram portion of this gel was calculated to contain 1000 mg of calcium carbonate. The organoleptic properties of this gel were similar to the product described in Example 1. After extended storage the product has shown no microbial growth and retains its desirable organoleptic properties.

15

Example 4

Chewable Aluminum Hydroxide/Magnesium Hydroxide Antacid Carrageenan-based
Confection

20 The method of Example 1 was followed except that 90 grams of magnesium hydroxide and 100 grams of aluminum hydroxide replaced the calcium carbonate and lemon flavoring replaced the wintergreen flavoring. Before pouring this formulation into the molds, it was determined that the solids content was 82.2% by weight. The gel was reported to be more sticky and paste-like than that of
25 Example 1. However, the gel had an easy bite and satisfactory flavor and mouthfeel. In addition, as the gel dissolved in the mouth, it left a feeling of fine sand remaining. A 5.02 gram portion of this gel was calculated to contain 450 mg of magnesium hydroxide and 500 mg of aluminum hydroxide.

Example 5

Chewable Acetaminophen Carrageenan-based Confection

5 The method of Example 1 was followed exactly except that ethyl cellulose-coated acetaminophen (108.7 grams) replaced the calcium carbonate. Before pouring this formulation into the molds, the solids content was determined to be 78.7% by weight. It was calculated that a 4.53 gram portion of the solidified gel contained 500 mg of acetaminophen. There were small white particles of
10 acetaminophen uniformly suspended in the gel; however, the sample had a smooth bite, and the taste of acetaminophen as well as its grittiness were eliminated.

Example 6

Chewable Ferrous Fumarate Carrageenan-based Confection

15 The method of Example 1 was followed except that 30.4 grams of ferrous fumarate replaced the calcium carbonate and spearmint flavor replaced the wintergreen flavor. In addition, 0.2% by weight of sodium benzoate was added to the formulation as a preservative. Before pouring this formulation into the molds,
20 the solids content was determined to be 78.3% by weight. It was calculated that a 4.06 gram portion of the solidified gel contained 49 milligrams of elemental iron. The formulation poured easily into the molds. The dark red gel was less sticky than previous experiments with iron supplements. When chewed, there was no toothpacking or sticking. Furthermore, there was no aftertaste of the iron that was
25 detectable.

Example 7

Chewable Multi-mineral Carrageenan-based Confection

30 The method of example 1 was followed except that the following mineral mixture was added to the syrup solution: 11.02 grams of ferrous fumarate, 111.12 grams of dicalcium phosphate \cdot 2H₂O, dibasic, 33.18 grams of magnesium oxide, 0.5 gram of cupric oxide, 3.73 grams of zinc oxide (40 mesh), 1.37 grams of

manganese sulfate, 0.039 gram of potassium iodide. As in Example 6, spearmint flavor and 0.2% by weight of sodium benzoate were added to the formulation. The resulting product chewed easily without any toothpacking or stickiness. However, there was some grittiness resulting from the use of granular dicalcium phosphate and the 40 mesh zinc oxide. Replacement of the granular dicalcium phosphate with powdered material was shown to eliminate much of the grittiness, and replacement of the 40 mesh zinc oxide with powdered zinc oxide would eliminate the remaining grittiness, making the product completely satisfactory.

10

Example 8

Chewable Calcium Carbonate/Ferrous Fumarate Carrageenan-based Confection

The method of Example 1 was followed except that 300 grams of calcium carbonate and 11.02 grams of ferrous fumarate were added to the syrup solution, and spearmint flavor and 0.2% by weight of sodium benzoate were used. The ferrous fumarate was added first and fully dispersed before the calcium carbonate was added in three equal portions. Each addition of calcium carbonate was followed by sufficient mixing to completely disperse it before adding the next portion. Before pouring this formulation, the solids content was calculated to be 79.1%. When chewed, the product had a clean bite without any iron aftertaste or a chalky taste. Also, there was no perceptible toothpacking or stickiness with this product. A 5.806 gram portion of the gelled product contained 1500 mg of calcium carbonate and 18 mg of elemental iron. While this invention has been described in terms of certain preferred embodiments and illustrated by specific examples, the invention is not to be construed as limited except as set forth in the following claims.

25

Claims

1. A chewable composition for delivery of a pharmacologically active material to a user comprising sweetener carrageenan and water, said composition
5 comprising from about 50% to about 83% of solids.
2. A chewable composition according to claim 1 wherein the hydrocolloid is carrageenan sulfated to the extent of from about 18% to about 35% based on the weight of the molecule.
10
3. A chewable composition according to claim 1 wherein the solids content of said composition comprises from about 78% to about 83% of solids and said chewable composition does not support microbial growth thereon.
4. A chewable composition according to claim 1 wherein said composition
15 is stable at acid, neutral and alkaline conditions.
5. A chewable composition according to claim 1 wherein said composition is stable at a pH of from about 2 to a pH of about 11.
20
6. A chewable composition according to claim 1 wherein carrageenan is present in an amount from about 2% to about 5.5% and optionally one or more of an additional hydrocolloid is present in a total amount of from about 0.5% to about 2%.
7. A chewable composition according to claim 6 wherein the additional
25 hydrocolloid is locust bean gum.
8. A chewable composition according to claim 6 wherein the additional hydrocolloid is a polysaccharide sulfated to an extent greater than 35% based on
30 the weight of the molecule.

9. A chewable composition according to claim 6 wherein the additional hydrocolloid is konjac mannan.

10. A chewable composition according to claim 6 wherein the additional hydrocolloid is a starch.

11. A chewable composition according to claim 6 wherein the additional hydrocolloid is guar gum.

12. A chewable composition according to claim 1 comprising a pharmacologically active material.

13. A chewable composition according to claim 12 wherein the pharmacologically active material is an antacid.

14. A chewable composition according to claim 12 wherein the pharmacologically active material is an antihistamine material.

15. A chewable composition according to claim 12 wherein the pharmacologically active material is an anti-inflammatory material or antipyretic material.

16. A chewable composition according to claim 12 wherein the pharmacologically active material is a nutritional supplement.

17. A chewable composition according to claim 12 wherein the pharmacologically active material is an antibiotic or antiviral material.

18. A chewable composition according to claim 12 wherein the pharmacologically active material is an expectorant or anti-tussive material.

19. A chewable composition according to claim 1 wherein the sweetener comprises sucrose.

20. A chewable composition according to claim 1 wherein the sweetener
5 comprises fructose.

21. A chewable composition according to claim 1 wherein the sweetener comprises galactose.

10 22. A chewable composition according to claim 1 wherein the sweetener is an artificial sweetener.

23. A chewable composition according to claim 22 wherein the sweetener is
selected from group consisting of aspartame, saccharin or cyclamate and mixtures
15 thereof.

24. A chewable composition according to claim 1 wherein an anionic or neutral preservative is present.

20 25. A chewable composition according to claim 24 wherein said preservative is present in an amount of from about 0.02% to about 0.5%.

26. A chewable composition according to claim 24 wherein said preservative is sodium benzoate, potassium sorbate, or a paraben.

25

27. A chewable composition according to claim 24 wherein said delivery system comprises from about 50% to up to about 78% of solids.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 9/20, A23G 3/00	A3	(11) International Publication Number: WO 98/20860 (43) International Publication Date: 22 May 1998 (22.05.98)
(21) International Application Number: PCT/US97/20217 (22) International Filing Date: 13 November 1997 (13.11.97) (30) Priority Data: 08/746,733 15 November 1996 (15.11.96) US (71) Applicant: FMC CORPORATION [US/US]; 1735 Market Street, Philadelphia, PA 19103 (US). (72) Inventors: BUBNIS, William, A.; 5904 Ravens Crest Drive, Plainsboro, NJ 08536 (US). REILLY, William, J., Jr.; 409 Byerly Drive, New Hope, PA 18938 (US). (74) Agent: CALDWELL, John; Woodcock Washburn Kurtz Mackiewicz & Norris, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 20 August 1998 (20.08.98)	
(54) Title: HYDROCOLLOID DELIVERY SYSTEM (57) Abstract A chewable delivery system stable in acidic, neutral or alkaline systems and having good mouthfeel is disclosed.		

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/20217

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K9/20 A23G3/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K A23G

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 379 147 A (STERLING DRUG INC) 25 July 1990 see page 4, line 1 - line 23 ---	1-18
A	US 5 298 261 A (PEBLEY WALTER S ET AL) 29 March 1994 see column 4, line 7 - line 24 see column 19; example 81 ---	1-18
A	DE 44 15 999 A (BOLDER ARZNEIMITTEL GMBH) 9 November 1995 see column 2, line 49 - column 3, line 45 ---	1-18
A	GB 2 160 771 A (LION CORP) 2 January 1986 see page 3; example 4 ---	1-18
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

3 July 1998

Date of mailing of the international search report

10/07/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Boulois, D

INTERNATIONAL SEARCH REPORT

Information on patent family members

In: International Application No

PCT/US 97/20217

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0379147	A	25-07-1990	US 5288479 A	22-02-1994
			AU 622259 B	02-04-1992
			AU 4786090 A	26-07-1990
			CA 2007752 A	17-07-1990
			DE 69012842 D	03-11-1994
			DE 69012842 T	20-04-1995
			DK 379147 T	12-12-1994
			FI 900099 A	18-07-1990
			IE 64163 B	12-07-1995
			JP 2247116 A	02-10-1990
			NO 180282 B	16-12-1996
US 5298261	A	29-03-1994	AU 5960294 A	19-07-1994
			WO 9414422 A	07-07-1994
DE 4415999	A	09-11-1995	AT 164061 T	15-04-1998
			DE 59501652 D	23-04-1998
			WO 9530407 A	16-11-1995
			EP 0758225 A	19-02-1997
			JP 9512800 T	22-12-1997
GB 2160771	A	02-01-1986	JP 61012629 A	21-01-1986
			CA 1275380 A	23-10-1990
			DE 3522898 A	02-01-1986
			US 4714612 A	22-12-1987
WO 9324103	A	09-12-1993	AU 4315793 A	30-12-1993
			EP 0641190 A	08-03-1995
			US 5688492 A	18-11-1997